

Molina Healthcare of Washington Medicaid Private Duty Nursing Prior Authorization Request Form Phone Number: (800) 869-7175

Fax Number: (800) 767-7188

		ı	MEMBE	ER INFORM	MATION							
Plan:	☐ Molina Medicaid (If Molina is secondary, please include a copy of the denial from primary insurance)											
Member Name:							DOB:		/		/	
Member ID#:							Phone:	()		-	
Service Type:	□ Electi	ve/Rou	ıtine 🗆	Expedited/	Urgent							
REFERRAL/SERVICE TYPE REQUESTED												
Diagnosis Code & D	escriptio)	n:										
CPT/HCPC Code & Description:												
90 DOS SPAN ONLY For continuation requests, the start date is always the day after the last authorization ends						DOS From: / / to / /						
		Р	ROVID	ER INFOR	MATIO	1						
Requesting Provide	· Name:				NPI#:			IN#:				
Servicing Provider or	g Provider or Facility:			NPI#:	TIN#:							
Contact at Rec Provider's									'			
Phone N	umber: (()	_		Num	Fax ber:	()		_			
CLINICAL DOC	UMENTA	TIONT	O SUPP	PORT NEED	FOR PRI	/ATE	DUTY N	UR	SING	(PE	N)	
Signed and dated physician order [Please submit: Home Health Certification Department of Health and Human S 0938-0357]			ication and Plan of Care,			□ Submitted						
Current history and (recent hospital adr Current treatment Current nursing control Recent daily nursing Emergency medical Plan and need for lifted places described.	missions/onissio	d treat - Most ng char	ment rerecent r	ecords notes (2 wee			ubmitted ubmitted ubmitted ubmitted ubmitted ubmitted		□ No	ot Si ot Si ot Si ot Si	ubmit ubmit ubmit ubmit ubmit	tted tted tted tted
	more tha	_	•	to supply co	ıre at a ti						JUIIII	<u> </u>

CLINICAL PRESENTATION (check all that apply)
Frequency of assessments (to include vital signs, interventions to support patient care, health
status assessment, etc.):
□ Once per 8 hour shift
□ 2-3 times per 8 hour shift
☐ Hourly or more often
Behavioral health, cognition, developmental monitoring:
□ Non-verbal, infrequent speech, or difficult to understand
☐ Self-abusive behavior, risk of self-harm, and intervention required
☐ Sleep disturbance and patient awake more than 3 hours per night
☐ Combative, confused, or disoriented behavior that impacts self-management; patient obese
☐ Combative, confused, or disoriented behavior that impacts self-management
Respiratory:
BiPAP/CPAP management
☐ More than 8 hours per day
☐ Less than 8 hours per day ☐ Nebulizer therapy
☐ More frequent than every 4 hours
☐ Every 4-24 hours
☐ Less frequent than daily, but at least once every 7 days
☐ Chest Physiotherapy – percussion, high-frequency chest wall oscillation vest, cough assist
device, etc.
☐ More than once per hour
☐ Every 1-4 hours
☐ Less than every 4 hours, but at least daily
□ Oxygen management
☐ Oxygen humidification, tracheal, no ventilator
☐ Oxygen needed at least weekly, based on pulse oximetry
☐ Suctioning
☐ Tracheal suctioning at least once every 2 hours
☐ Tracheal suctioning daily, but less than every 2 hours
☐ Nasal or oral suctioning daily
☐ Tracheostomy management
☐ Tracheostomy management with complications (skin breakdown, replacement needed)
☐ Tracheostomy management, no complications
□ Ventilator management□ Continuous ventilator use
☐ Ventilator use for 12 or more hours per day
☐ Ventilator use for 7-12 hours per day
☐ Ventilator use for less than 7 hours per day
☐ Interventions in place for active weaning
☐ Ventilator weaning achieved; requires ongoing post-weaning monitoring and management
☐ Ventilator on standby, respiratory assistance, or used at night for less than 1 hour

Skilled Nursing Needs:	
☐ Blood draw	
	Peripheral line
·	Less than twice per week
☐ Infusion therapy	
☐ Blood or blood product	
☐ Chemotherapy infusion	
☐ Central line access and management	
☐ Pain medication infusion	
☐ Intravenous Infusion (IV antibiotics, etc.), including in	fusion administration and monitoring
for infusion reactions	
☐ Infusions more than every 4 hours	
☐ Infusions less than every 4 hours	
☐ Non-infusion medication	
☐ Insulin administration	
☐ Non-insulin medication injectable administra	tion
☐ Medication administration at least every 2 ho	urs, requiring clinical monitoring
☐ Activity of Daily Living (ADL)/Therapy support	
BedboundWhe	elchair user Ambulatory
☐ Total/partial lift, weight 55-125 pounds	
☐ Total/partial lift, weight greater than 125 poun	ds
☐ ADL support needed more than 4 hours per d	ay to maximize patient's independence
☐ Body cast management	
☐ Cast or brace management	
 Splinting management, including removal and 	
☐ Communication deficit; nurse to support there	
☐ Range of motion exercises at least every 8 ho	
☐ Physical therapy program at least 3 hours pe	er day; occupational therapy program
at least 4 hours per day	
☐ Nutrition management	
☐ Enteral nutrition with complications, requires o	9
adjustment or placement of tube, and assessr	ment or management of complications
 Enteral nutrition without complications 	
☐ Gastrostomy tube care, uncomplicated	
☐ Nasogastric tube care, uncomplicated	
☐ Partial parenteral nutrition with central line ca	
☐ Total parenteral nutrition with central line care	
☐ Skin and wound care management	
☐ Burn care	
☐ Ostomy care, at least once per day	
☐ Postsurgical care, within 45 days of surgery	
☐ Stage 1 or 2 wound management, at least one	
☐ Stage 3 or 4 wound management, at least on	•
☐ Stage 3 or 4 wound management at least one	
☐ Prescribed topical medication application at I	east every 4 hours
☐ Wound vacuum management	

□ Seizure control that requires nursing intervention/management □ Seizures lasting less than 3 minutes, at least 4 times per week □ Seizures lasting 3-5 minutes, at least 4 times per week □ Seizures lasting 3-5 minutes, 1 to 4 times per day □ Seizures lasting 3-5 minutes, more than 5 times per day □ Seizures lasting more than 5 minutes, or clustered seizures, or seizure activity without regaining consciousness, at least 4 times per week □ Seizures lasting more than 5 minutes, or clustered seizures, or seizure activity without regaining consciousness, one time or more per day, requiring rectal medication □ Seizures lasting more than 5 minutes, or clustered seizures, or seizure activity without regaining consciousness, one time or more per day, requiring IM or IV medication
ADDITIONAL INFORMATION
List:

MHW PART #1767-2201 MHW-2/23/2022 27916FRMMDWAEN 220225